

# Root hairs increase rhizosphere extension and carbon input to soil

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

---

## Abstract

© The Author(s) 2017. Published by Oxford University Press on behalf of the Annals of Botany Company. All rights reserved. Background and Aims Although it is commonly accepted that root exudation enhances plant-microbial interactions in the rhizosphere, experimental data on the spatial distribution of exudates are scarce. Our hypothesis was that root hairs exude organic substances to enlarge the rhizosphere farther from the root surface. Methods Barley (*Hordeum vulgare* 'Pallas' - wild type) and its root-hairless mutant (brb) were grown in rhizoboxes and labelled with  $^{14}\text{C}$ . A filter paper was placed on the soil surface to capture, image and quantify root exudates. Key Results Plants with root hairs allocated more carbon (C) to roots (wild type: 13 %; brb: 8 % of assimilated  $^{14}\text{C}$ ) and to rhizosheaths (wild type: 1.2 %; brb: 0.2 %), while hairless plants allocated more C to shoots (wild type: 65 %; brb: 75 %). Root hairs increased the radial rhizosphere extension three-fold, from 0.5 to 1.5 mm. Total exudation on filter paper was three times greater for wild type plants compared to the hairless mutant. Conclusion Root hairs increase exudation and spatial rhizosphere extension, which probably enhance rhizosphere interactions and nutrient cycling in larger soil volumes. Root hairs may therefore be beneficial to plants under nutrient-limiting conditions. The greater C allocation below ground in the presence of root hairs may additionally foster C sequestration.

<http://dx.doi.org/10.1093/aob/mcx127>

---

## Keywords

$^{14}\text{C}$  imaging, Barley (*Hordeum vulgare* L.), Carbon allocation, Rhizosphere extension, Root exudates, Root hairs, Root-soil interface

## References

- [1] Bardgett RD, Mommer L, De Vries FT. 2014. Going underground: root traits as drivers of ecosystem processes. *Trends in Ecology & Evolution* 29: 692-699.
- [2] Bates TR, Lynch JP. 1996. Stimulation of root hair elongation in *Arabidopsis thaliana* by low phosphorus availability. *Plant, Cell & Environment* 19: 529-538.
- [3] Bertin C, Yang X, Weston LA. 2003. The role of root exudates and allelochemicals in the rhizosphere. *Plant and Soil* 256: 67-83.
- [4] Brown LK, George TS, Thompson JA, et al. 2012. What are the implications of variation in root hair length on tolerance to phosphorus deficiency in combination with water stress in barley (*Hordeum vulgare*)? *Annals of Botany* 110: 319-328.
- [5] Czarnota MA, Paul RN, Weston LA, Duke SO. 2003. Anatomy of sorgoleone-secreting root hairs of *Sorghum* species. *International Journal of Plant Sciences* 164: 861-866.

- [6] Dakora FD, Phillips DA. 2002. Root exudates as mediators of mineral acquisition in low-nutrient environments. *Plant and Soil* 245: 35-47.
- [7] Dart PJ. 1971. Scanning electron microscopy of plant roots. *Journal of Experimental Botany* 22: 163-168.
- [8] Dayan FE, Howell J, Weidenhamer JD. 2009. Dynamic root exudation of sorgoleone and its in planta mechanism of action. *Journal of Experimental Botany* 60: 2107-2177.
- [9] Dennis PG, Miller AJ, Hirsch PR. 2010. Are root exudates more important than other sources of rhizodeposits in structuring rhizosphere bacterial communities? *FEMS Microbiology Ecology* 72: 313-327.
- [10] FAO. 2012. Coping with water scarcity: An action framework for agriculture and food security. Water Report 38. Rome: Food and Agriculture Organization.
- [11] Gahoonia TS, Care D, Nielsen NE. 1997. Root hairs and phosphorus acquisition of wheat and barley cultivars. *Plant and Soil* 191: 181-188.
- [12] Gilroy S, Jones DL. 2000. Through form to function: root hair development and nutrient uptake. *Trends in Plant Sciences* 5: 1360-1385.
- [13] Greaves MP, Darbyshire JF. 1972. The ultrastructure of the mucilaginous layer on plant roots. *Soil Biology & Biochemistry* 4: 443-449.
- [14] Gunina A, Kuzyakov Y. 2015. Sugars in soil and sweets for microorganisms: review of origin, content, composition and fate. *Soil Biology and Biochemistry* 90: 87-100.
- [15] Haling RE, Simpson RJ, Delhaize E, Hocking PJ, Richardson AE. 2010. Effect of lime on root growth, morphology and the rhizosphere of cereal seedlings growing in an acid soil. *Plant and Soil* 327: 199-212.
- [16] Hamilton EW, Frank DA. 2001. Can plants stimulate soil microbes and their own nutrient supply? Evidence from a grazing tolerant grass. *Ecology* 82: 2397-2402.
- [17] Herman DJ, Johnson KK, Jaeger CH, Schwartz E, Firestone MK. 2006. Root influence on nitrogen mineralization and nitrification in rhizosphere soil. *Soil Science Society of America Journal* 70: 1504.
- [18] Hinsinger P. 2001. Bioavailability of soil inorganic P in the rhizosphere as affected by root-induced chemical changes: a review. *Plant and Soil* 237: 173-195.
- [19] Jones DL, Nguyen C, Finlay RD. 2009. Carbon flow in the rhizosphere: carbon trading at the soil-root interface. *Plant and Soil* 321: 5-33.
- [20] Jungk A. 2001. Root hairs and the acquisition of plant nutrients from soil. *Journal of Plant Nutrition and Soil Science* 164: 121-129.
- [21] Kuzyakov Y, Xu X. 2013. Tansley review: Competition between roots and microorganisms for nitrogen: mechanisms and ecological relevance. *New Phytologist* 198: 656-669.
- [22] Kuzyakov Y, Shevtzova E, Pustovoytov K. 2006. Carbonate re-crystallization in soil revealed by <sup>14</sup>C labeling: experiment, model and significance for paleo-environmental reconstructions. *Geoderma* 131: 45-58.
- [23] Landi L, Valori F, Ascher J, Renella G, Falchini L, Nannipieri P. 2006. Root exudate effects on the bacterial communities, CO<sub>2</sub> evolution, nitrogen transformations and ATP content of rhizosphere and bulk soils. *Soil Biology and Biochemistry* 38: 509-516.
- [24] Leinweber P, Eckhardt KU, Fischer H, Kuzyakov Y. 2008. A new rapid micro-method for the molecular-chemical characterization of rhizodeposits by field-ionization mass spectrometry. *Rapid Communications in Mass Spectrometry* 22: 1230-1234.
- [25] Lobet G, Pagès L, Draye X. 2011. A novel image-analysis toolbox enabling quantitative analysis of root system architecture. *Plant Physiology* 157: 29-39.
- [26] Lynch JP. 2007. Roots of the second green revolution. *Australian Journal of Botany* 55: 493-512.
- [27] Marschner P, Crowley D, Rengel Z. 2011. Rhizosphere interactions between microorganisms and plants govern iron and phosphorus acquisition along the root axis-model and research methods. *Soil Biology and Biochemistry* 43: 883-894.
- [28] Parry MAJ, Hawkesford MJ. 2010. Food security: increasing yield and improving resource use efficiency. *Proceedings of the Nutrition Society* 69: 592-600.
- [29] Parry MAJ, Flexas J, Medrano H. 2005. Prospects for crop production under drought: research priorities and future directions. *Annals of Applied Biology* 147: 211-226.
- [30] Pausch J, Loepmann S, Kühnel A, Forbush K. 2016. Rhizosphere priming of barley with and without root hairs. *Soil Biology & Biochemistry* 100: 74-82.
- [31] Peterson RL, Farquhar ML. 1996. Root hairs: specialized tubular cells extending root surfaces. *Botanical Review* 62: 1-40.
- [32] Rasse DP, Rumpel C, Dignac MF. 2005. Is soil carbon mostly root carbon? Mechanisms for a specific stabilisation. *Plant and Soil* 269: 341-356.
- [33] Sanaullah M, Chabbi A, Rumpel C, Kuzyakov Y. 2012. Carbon allocation in grassland communities under drought stress followed by <sup>14</sup>C pulse labeling. *Soil Biology and Biochemistry* 55: 132-139.

- [34] Schweiger PF, Robson AD, Barrow NJ. 1995. Root hair length determines beneficial effect of a *Glomus* species on shoot growth of some pasture species. *New Phytologist* 131: 247-254.
- [35] Segal E, Kushnir T, Mualem Y, Shani U. 2008. Water uptake and hydraulics of the root hair rhizosphere. *Vadose Zone Journal* 7: 1027-1034.
- [36] Sprent JI. 1975. Adherence of sand particles to soybean roots under water stress. *New Phytologist* 74: 461-463.
- [37] Weston LA, Ryan PR, Watt M. 2012. Mechanisms for cellular transport and release of allelochemicals from plant roots into the rhizosphere. *Journal of Experimental Botany* 63: 3445-3454.
- [38] Yan X, Liao H, Beebe SE, Blair MW, Lynch JP. 2004. QTL mapping of root hair and acid exudation traits and their relationship to phosphorus uptake in common bean. *Plant and Soil* 265: 17-29.
- [39] York LM, Carminati A, Mooney SJ, Ritz K, Bennett MJ. 2016. The holistic rhizosphere: integrating zones, processes, and semantics in the soil influenced by roots. *Journal of Experimental Botany* 67: 3629-3643.